آگهی سخنرانی

The Sphere Covering Inequality and Its Applications

امير مرادي فام

University of California, Riverside, USA

Abstarct:

We show that the total area of two distinct Gaussian curvature 1 surfaces with the same conformal factor on the boundary, which are also conformal to the Euclidean unit disk, must be at least 4π . In other words, the areas of these surfaces must cover the whole unit sphere after a proper rearrangement. We refer to this lower bound of total areas as the Sphere Covering Inequality. This inequality and its generalizations are applied to a number of open problems related to Moser-Trudinger type inequalities, mean field equations and Onsager vortices, etc, and yield optimal results. In particular we confirm the best constant of a Moser-Truidinger type inequality conjectured by A. Chang and P. Yang in 1987. This is a joint work with Changfeng Gui.

زمان:

شنبه ۱۶ مردادماه ۱۳۹۵ ساعت: ۱۱:۳۰ الی ۱۲:۴۵ یکشنبه ۱۷ مردادماه ۱۳۹۵ ساعت: ۱۱:۳۰ الی ۱۲:۴۵ یکشنبه ۱۷ مردادماه ۱۳۹۵ ساعت: ۱۴:۰۰ الی ۱۵:۱۵

مکان: میدان نیاوران، پژوهشگاه دانشهای بنیادی، پژوهشکده ریاضیات،

سال ا