

Alternating Permutations

Richard P. Stanley

M.I.T.

Definitions

A sequence a_1, a_2, \dots, a_k of distinct integers is **alternating** if

$$a_1 > a_2 < a_3 > a_4 < \cdots ,$$

and **reverse alternating** if

$$a_1 < a_2 > a_3 < a_4 > \cdots .$$

Definitions

A sequence a_1, a_2, \dots, a_k of distinct integers is **alternating** if

$$a_1 > a_2 < a_3 > a_4 < \cdots ,$$

and **reverse alternating** if

$$a_1 < a_2 > a_3 < a_4 > \cdots .$$

\mathfrak{S}_n : symmetric group of all permutations of
 $1, 2, \dots, n$

E_n

$$\begin{aligned}\textcolor{red}{E_n} &= \#\{w \in \mathfrak{S}_n : w \text{ is alternating}\} \\ &= \#\{w \in \mathfrak{S}_n : w \text{ is reverse alternating}\}\end{aligned}$$

via

$$a_1 a_2 \cdots a_n \mapsto n + 1 - a_1, n + 1 - a_2, \dots, n + 1 - a_n$$

E_n

$$\begin{aligned}\textcolor{red}{E_n} &= \#\{w \in \mathfrak{S}_n : w \text{ is alternating}\} \\ &= \#\{w \in \mathfrak{S}_n : w \text{ is reverse alternating}\}\end{aligned}$$

via

$$a_1 a_2 \cdots a_n \mapsto n + 1 - a_1, n + 1 - a_2, \dots, n + 1 - a_n$$

E.g., $E_4 = 5$: 2143, 3142, 3241, 4132, 4231

André's theorem

Theorem (Désiré André, 1879)

$$y := \sum_{n \geq 0} E_n \frac{x^n}{n!} = \sec x + \tan x$$

André's theorem

Theorem (Désiré André, 1879)

$$y := \sum_{n \geq 0} E_n \frac{x^n}{n!} = \sec x + \tan x$$

E_n is an **Euler number**,

E_{2n} a **secant number**,

E_{2n-1} a **tangent number**.

Naive proof

52439 **1** 826
 \leftarrow \rightarrow
alternating *alternating*

Naive proof



Obtain w that is **either** alternating or reverse alternating.

Naive proof

52439 **1** 826
 \leftarrow \rightarrow
alternating *alternating*

Obtain w that is **either** alternating or reverse alternating.

$$\Rightarrow 2E_{n+1} = \sum_{k=0}^n \binom{n}{k} E_k E_{n-k}, \quad n \geq 1$$

Completion of proof

$$2E_{n+1} = \sum_{k=0}^n \binom{n}{k} E_k E_{n-k}, \quad n \geq 1$$

Completion of proof

$$2E_{n+1} = \sum_{k=0}^n \binom{n}{k} E_k E_{n-k}, \quad n \geq 1$$

$$2 \sum_{n \geq 0} E_{n+1} \frac{x^n}{n!} = 2y'$$

$$\sum_{n \geq 0} \left(\sum_{k=0}^n \binom{n}{k} E_k E_{n-k} \right) \frac{x^n}{n!} = 1 + y^2$$

Completion of proof

$$2E_{n+1} = \sum_{k=0}^n \binom{n}{k} E_k E_{n-k}, \quad n \geq 1$$

$$2 \sum_{n \geq 0} E_{n+1} \frac{x^n}{n!} = 2y'$$

$$\sum_{n \geq 0} \left(\sum_{k=0}^n \binom{n}{k} E_k E_{n-k} \right) \frac{x^n}{n!} = 1 + y^2$$

$$\Rightarrow 2y' = 1 + y^2, \text{ etc.}$$

A generalization

\exists more sophisticated approaches.

E.g., let

$$f_k(n) = \#\{w \in \mathfrak{S}_n : w(r) < w(r+1) \Leftrightarrow k|r\}$$

$$f_2(n) = E_n$$

A generalization

\exists more sophisticated approaches.

E.g., let

$$f_k(n) = \#\{w \in \mathfrak{S}_n : w(r) < w(r+1) \Leftrightarrow k|r\}$$

$$f_2(n) = E_n$$

Theorem.

$$\sum_{k \geq 0} f_k(kn) \frac{x^{kn}}{(kn)!} = \left(\sum_{n \geq 0} (-1)^n \frac{x^{kn}}{(kn)!} \right)^{-1}.$$

Combinatorial trigonometry

Define $\sec x$ and $\tan x$ by

$$\sec x + \tan x = \sum_{n \geq 0} E_n \frac{x^n}{n!}.$$

Combinatorial trigonometry

Define $\sec x$ and $\tan x$ by

$$\sec x + \tan x = \sum_{n \geq 0} E_n \frac{x^n}{n!}.$$

Exercise. $\sec^2 x = 1 + \tan^2 x$

Combinatorial trigonometry

Define $\sec x$ and $\tan x$ by

$$\sec x + \tan x = \sum_{n \geq 0} E_n \frac{x^n}{n!}.$$


Exercise. $\sec^2 x = 1 + \tan^2 x$

Exercise. $\tan(x + y) = \frac{\tan x + \tan y}{1 - (\tan x)(\tan y)}$


Some occurrences of Euler numbers

(1) E_{2n-1} is the number of complete increasing binary trees on the vertex set $[2n + 1] = \{1, 2, \dots, 2n + 1\}$.

Five vertices



Five vertices



Slightly more complicated for E_{2n}

Simsun permutations

Definition. A permutation $w = a_1 \cdots a_n \in \mathfrak{S}_n$ is **simsun** if for all $1 \leq k \leq n$, the restriction $b_1 \cdots b_k$ of w to $1, 2, \dots, k$ has no **double descents** $b_{i-1} > b_i > b_{i+1}$.

Simsun permutations

Definition. A permutation $w = a_1 \cdots a_n \in \mathfrak{S}_n$ is **simsun** if for all $1 \leq k \leq n$, the restriction $b_1 \cdots b_k$ of w to $1, 2, \dots, k$ has no **double descents** $b_{i-1} > b_i > b_{i+1}$.

Example. 346251 is **not** simsun: restriction to 1, 2, 3, 4 is **3421**.

Simsun permutations

Definition. A permutation $w = a_1 \cdots a_n \in \mathfrak{S}_n$ is **simsun** if for all $1 \leq k \leq n$, the restriction $b_1 \cdots b_k$ of w to $1, 2, \dots, k$ has no **double descents** $b_{i-1} > b_i > b_{i+1}$.

Example. 346251 is **not** simsun: restriction to 1, 2, 3, 4 is **3421**.

Theorem (R. Simion and S. Sundaram) *The number of simsun permutations $w \in \mathfrak{S}_n$ is E_n .*

Chains of partitions

(2) Start with n one-element sets $\{1\}, \dots, \{n\}$.

Chains of partitions

- (2) Start with n one-element sets $\{1\}, \dots, \{n\}$.
Merge together two at a time until reaching
 $\{1, 2, \dots, n\}$.

Chains of partitions

(2) Start with n one-element sets $\{1\}, \dots, \{n\}$.

Merge together two at a time until reaching
 $\{1, 2, \dots, n\}$.

$1-2-3-4-5-6, \quad 12-3-4-5-6, \quad 12-34-5-6$

$125-34-6, \quad 125-346, \quad 123456$

Chains of partitions

(2) Start with n one-element sets $\{1\}, \dots, \{n\}$.

Merge together two at a time until reaching $\{1, 2, \dots, n\}$.

1–2–3–4–5–6, 12–3–4–5–6, 12–34–5–6

125–34–6, 125–346, 123456

S_n acts on these chains.

Chains of partitions

(2) Start with n one-element sets $\{1\}, \dots, \{n\}$.

Merge together two at a time until reaching $\{1, 2, \dots, n\}$.

$1-2-3-4-5-6, \quad 12-3-4-5-6, \quad 12-34-5-6$

$125-34-6, \quad 125-346, \quad 123456$

\mathfrak{S}_n acts on these chains.

Theorem. *The number of \mathfrak{S}_n -orbits is E_{n-1} .*

Orbit representatives for $n = 5$

12–3–4–5

123–4–5

1234–5

12–3–4–5

123–4–5

123–45

12–3–4–5

12–34–5

125–34

12–3–4–5

12–34–5

12–345

12–3–4–5

12–34–5

1234–5

A polytope volume

Let \mathcal{P}_n be the convex polytope in \mathbb{R}^n with equations

$$x_i \geq 0, \quad 1 \leq i \leq n$$

$$x_i + x_{i+1} \leq 1, \quad 1 \leq i \leq n - 1.$$

A polytope volume

Let \mathcal{P}_n be the convex polytope in \mathbb{R}^n with equations

$$\begin{aligned}x_i &\geq 0, \quad 1 \leq i \leq n \\x_i + x_{i+1} &\leq 1, \quad 1 \leq i \leq n - 1.\end{aligned}$$

Theorem. $\text{vol}(\mathcal{P}_n) = \frac{1}{n!} E_n$

Ulam's problem

Let $E(n)$ be the expected length $\text{is}(w)$ of the longest **increasing** subsequence of w .

Ulam's problem

Let $E(n)$ be the expected length $\text{is}(w)$ of the longest **increasing** subsequence of w .

$$w = 5\mathbf{2}41\mathbf{7}36 \Rightarrow \text{is}(w) = 3$$

Two highlights

Long story ...

Two highlights

Long story ...

Logan-Shepp, Vershik-Kerov:

$$E(n) \sim 2\sqrt{n}$$

Two highlights

Long story ...

Logan-Shepp, Vershik-Kerov:

$$E(n) \sim 2\sqrt{n}$$

Write $\text{is}_n(w)$ for $\text{is}(w)$ when $w \in \mathfrak{S}_n$.

Baik-Deift-Johansson:

$$\lim_{n \rightarrow \infty} \text{Prob} \left(\frac{\text{is}_n(w) - 2\sqrt{n}}{n^{1/6}} \leq t \right) = F(t),$$

the **Tracy-Widom distribution**.

Alternating subsequences?

as(w) = length of longest alternating subseq. of w

$$w = \mathbf{5}6\mathbf{2}1\mathbf{8}3\mathbf{4}7 \Rightarrow \text{as}(w) = 5$$

The main lemma

MAIN LEMMA. $\forall w \in \mathfrak{S}_n \exists$ alternating subsequence of maximal length that contains n .

The main lemma

MAIN LEMMA. $\forall w \in \mathfrak{S}_n \exists$ alternating subsequence of maximal length that contains n .

$$a_k(n) = \#\{w \in \mathfrak{S}_n : \text{as}(w) = k\}$$

$$\begin{aligned} b_k(n) &= a_1(n) + a_2(n) + \cdots + a_k(n) \\ &= \#\{w \in \mathfrak{S}_n : \text{as}(w) \leq k\}. \end{aligned}$$

Recurrence for $a_k(n)$

$$\Rightarrow a_k(n) = \sum_{j=1}^n \binom{n-1}{j-1}$$

$$\sum_{2r+s=k-1} (a_{2r}(j-1) + a_{2r+1}(j-1)) a_s(n-j)$$

The main generating function

Define $B(x, t) = \sum_{k,n \geq 0} b_k(n) t^k \frac{x^n}{n!}$

Theorem.

$$B(x, t) = \frac{2/\rho}{1 - \frac{1-\rho}{t} e^{\rho x}} - \frac{1}{\rho},$$

where $\rho = \sqrt{1 - t^2}$.

Formulas for $b_k(n)$

Corollary.

$$\Rightarrow b_1(n) = 1$$

$$b_2(n) = n$$

$$b_3(n) = \frac{1}{4}(3^n - 2n + 3)$$

$$b_4(n) = \frac{1}{8}(4^n - (2n - 4)2^n)$$

⋮

Formulas for $b_k(n)$

Corollary.

$$\Rightarrow b_1(n) = 1$$

$$b_2(n) = n$$

$$b_3(n) = \frac{1}{4}(3^n - 2n + 3)$$

$$b_4(n) = \frac{1}{8}(4^n - (2n - 4)2^n)$$

⋮

no such formulas for longest **increasing** subsequences

Mean (expectation) of $\text{as}(w)$

$$D(n) = \frac{1}{n!} \sum_{w \in \mathfrak{S}_n} \text{as}(w),$$

the **expectation** of $\text{as}(n)$ for $w \in \mathfrak{S}_n$

Mean (expectation) of $\text{as}(w)$

$$D(n) = \frac{1}{n!} \sum_{w \in \mathfrak{S}_n} \text{as}(w),$$

the **expectation** of $\text{as}(n)$ for $w \in \mathfrak{S}_n$

Let

$$A(x, t) = \sum_{k, n \geq 0} a_k(n) t^k \frac{x^n}{n!} = (1 - t) B(x, t)$$

$$= (1 - t) \left(\frac{2/\rho}{1 - \frac{1-\rho}{t} e^{\rho x}} - \frac{1}{\rho} \right).$$

Formula for $D(n)$

$$\begin{aligned}\sum_{n \geq 0} D(n)x^n &= \frac{\partial}{\partial t} A(x, 1) \\&= \frac{6x - 3x^2 + x^3}{6(1-x)^2} \\&= x + \sum_{n \geq 2} \frac{4n+1}{6} x^n.\end{aligned}$$

Formula for $D(n)$

$$\begin{aligned}\sum_{n \geq 0} D(n)x^n &= \frac{\partial}{\partial t} A(x, 1) \\&= \frac{6x - 3x^2 + x^3}{6(1-x)^2} \\&= x + \sum_{n \geq 2} \frac{4n+1}{6} x^n.\end{aligned}$$
$$\Rightarrow A(n) = \frac{4n+1}{6}, \quad n \geq 2$$

Formula for $D(n)$

$$\begin{aligned}\sum_{n \geq 0} D(n)x^n &= \frac{\partial}{\partial t} A(x, 1) \\&= \frac{6x - 3x^2 + x^3}{6(1-x)^2} \\&= x + \sum_{n \geq 2} \frac{4n+1}{6} x^n.\end{aligned}$$
$$\Rightarrow A(n) = \frac{4n+1}{6}, \quad n \geq 2$$

Recall $E(n) \sim 2\sqrt{n}$.

Variance of $\text{as}(w)$

$$V(n) = \frac{1}{n!} \sum_{w \in \mathfrak{S}_n} \left(\text{as}(w) - \frac{4n+1}{6} \right)^2, \quad n \geq 2$$

the **variance** of $\text{as}(n)$ for $w \in \mathfrak{S}_n$

Variance of $\text{as}(w)$

$$V(n) = \frac{1}{n!} \sum_{w \in \mathfrak{S}_n} \left(\text{as}(w) - \frac{4n+1}{6} \right)^2, \quad n \geq 2$$

the **variance** of $\text{as}(n)$ for $w \in \mathfrak{S}_n$

Corollary.

$$V(n) = \frac{8}{45}n - \frac{13}{180}, \quad n \geq 4$$

Variance of $\text{as}(w)$

$$V(n) = \frac{1}{n!} \sum_{w \in \mathfrak{S}_n} \left(\text{as}(w) - \frac{4n+1}{6} \right)^2, \quad n \geq 2$$

the **variance** of $\text{as}(n)$ for $w \in \mathfrak{S}_n$

Corollary.

$$V(n) = \frac{8}{45}n - \frac{13}{180}, \quad n \geq 4$$

similar results for higher moments

A new distribution?

$$\mathbf{P}(t) = \lim_{n \rightarrow \infty} \text{Prob}_{w \in \mathfrak{S}_n} \left(\frac{\text{as}(w) - 2n/3}{\sqrt{n}} \leq t \right)$$

A new distribution?

$$P(t) = \lim_{n \rightarrow \infty} \text{Prob}_{w \in \mathfrak{S}_n} \left(\frac{\text{as}(w) - 2n/3}{\sqrt{n}} \leq t \right)$$

Stanley distribution?

Limiting distribution

Theorem (Pemantle, Widom, (Wilf)).

$$\lim_{n \rightarrow \infty} \text{Prob}_{w \in \mathfrak{S}_n} \left(\frac{\text{as}(w) - 2n/3}{\sqrt{n}} \leq t \right)$$

$$= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{t\sqrt{45}/4} e^{-s^2} ds$$

(Gaussian distribution)

Limiting distribution

Theorem (Pemantle, Widom, (Wilf)).

$$\lim_{n \rightarrow \infty} \text{Prob}_{w \in \mathfrak{S}_n} \left(\frac{\text{as}(w) - 2n/3}{\sqrt{n}} \leq t \right)$$

$$= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{t\sqrt{45}/4} e^{-s^2} ds$$

(Gaussian distribution)



Umbral enumeration

Umbral formula: involves E^k , where E is an indeterminate (the **umbra**). Replace E^k with the Euler number E_k . (Technique from 19th century, modernized by Rota et al.)

Umbral enumeration

Umbral formula: involves E^k , where E is an indeterminate (the **umbra**). Replace E^k with the Euler number E_k . (Technique from 19th century, modernized by Rota et al.)

Example.

$$\begin{aligned}(1 + E^2)^3 &= 1 + 3E^2 + 3E^4 + E^6 \\&= 1 + 3E_2 + 3E_4 + E_6 \\&= 1 + 3 \cdot 1 + 3 \cdot 5 + 61 \\&= 80\end{aligned}$$

Another example

$$\begin{aligned}(1+t)^E &= 1 + Et + \binom{E}{2}t^2 + \binom{E}{3}t^3 + \dots \\&= 1 + Et + \frac{1}{2}E(E-1)t^2 + \dots \\&= 1 + E_1t + \frac{1}{2}(E_2 - E_1))t^2 + \dots \\&= 1 + t + \frac{1}{2}(1-1)t^2 + \dots \\&= 1 + t + O(t^3).\end{aligned}$$

Alt. fixed-point free involutions

fixed point free involution $w \in \mathfrak{S}_{2n}$: all cycles of length two

Alt. fixed-point free involutions

fixed point free involution $w \in \mathfrak{S}_{2n}$: all cycles of length two

Let $f(n)$ be the number of **alternating** fixed-point free involutions in \mathfrak{S}_{2n} .

Alt. fixed-point free involutions

fixed point free involution $w \in \mathfrak{S}_{2n}$: all cycles of length two

Let $f(n)$ be the number of **alternating** fixed-point free involutions in \mathfrak{S}_{2n} .

$$n = 3 : \begin{aligned} 214365 &= (1, 2)(3, 4)(5, 6) \\ 645231 &= (1, 6)(2, 4)(3, 5) \end{aligned}$$

$$f(3) = 2$$

An umbral theorem

Theorem.

$$\mathbf{F}(x) = \sum_{n \geq 0} f(n)x^n$$

An umbral theorem

Theorem.

$$\begin{aligned} \mathbf{F}(x) &= \sum_{n \geq 0} f(n)x^n \\ &= \left(\frac{1+x}{1-x} \right)^{(E^2+1)/4} \end{aligned}$$

An umbral theorem

Theorem.

$$\begin{aligned} \mathbf{F}(x) &= \sum_{n \geq 0} f(n)x^n \\ &= \left(\frac{1+x}{1-x} \right)^{(E^2+1)/4} \end{aligned}$$

Proof. Uses representation theory of the symmetric group \mathfrak{S}_n . In particular, there is a “nice” representation of dimension E_n .

Ramanujan's Lost Notebook

Theorem (Ramanujan, Berndt, implicitly) As $x \rightarrow 0+$,

$$2 \sum_{n \geq 0} \left(\frac{1-x}{1+x} \right)^{n(n+1)} \sim \sum_{k \geq 0} f(k)x^k = F(x),$$

an **analytic** (non-formal) identity.

A formal identity

Corollary (via Ramanujan, Andrews).

$$F(x) = 2 \sum_{n \geq 0} q^n \frac{\prod_{j=1}^n (1 - q^{2j-1})}{\prod_{j=1}^{2n+1} (1 + q^j)},$$

where $\mathbf{q} = \left(\frac{1-x}{1+x}\right)^{2/3}$, a **formal identity**.

Simple result, hard proof

Recall: number of n -cycles in \mathfrak{S}_n is $(n - 1)!$.

Simple result, hard proof

Recall: number of n -cycles in \mathfrak{S}_n is $(n - 1)!$.

Theorem. Let $b(n)$ be the number of *alternating* n -cycles in \mathfrak{S}_n . Then if n is odd,

$$b(n) = \frac{1}{n} \sum_{d|n} \mu(d) (-1)^{(d-1)/2} E_{n/d}.$$

Special case

Corollary. *Let p be an odd prime. Then*

$$b(p) = \frac{1}{p} \left(E_p - (-1)^{(p-1)/2} \right).$$

Special case

Corollary. *Let p be an odd prime. Then*

$$b(p) = \frac{1}{p} \left(E_p - (-1)^{(p-1)/2} \right).$$

Combinatorial proof?

Another such result

Recall: the expected number of fixed points of $w \in \mathfrak{S}_n$ is 1.

Another such result

Recall: the expected number of fixed points of $w \in \mathfrak{S}_n$ is 1.

$J(n)$ = expected number of fixed points of
reverse alternating $w \in \mathfrak{S}_n$

Theorem. $J(2n) = \frac{1}{E_{2n}} (E_{2n} - (-1)^n).$

Similar (but more complicated) for n odd or for alternating permutations.

Another such result

Recall: the expected number of fixed points of $w \in \mathfrak{S}_n$ is 1.

$J(n)$ = expected number of fixed points of
reverse alternating $w \in \mathfrak{S}_n$

Theorem. $J(2n) = \frac{1}{E_{2n}} (E_{2n} - (-1)^n).$

Similar (but more complicated) for n odd or for alternating permutations.

Combinatorial proof?

Descent sets

Let $w = a_1 a_2 \cdots a_n \in \mathfrak{S}_n$. **Descent set** of w :

$$D(w) = \{i : a_i > a_{i+1}\} \subseteq \{1, \dots, n-1\}$$

Descent sets

Let $w = a_1 a_2 \cdots a_n \in \mathfrak{S}_n$. **Descent set** of w :

$$D(w) = \{i : a_i > a_{i+1}\} \subseteq \{1, \dots, n-1\}$$

$$D(4157623) = \{1, 4, 5\}$$

$$D(4152736) = \{1, 3, 5\} \text{ (alternating)}$$

$$D(4736152) = \{2, 4, 6\} \text{ (reverse alternating)}$$

$\beta_n(S)$

$$\beta_n(S) = \#\{w \in \mathfrak{S}_n : D(w) = S\}$$

$\beta_n(S)$

$$\beta_n(S) = \#\{w \in \mathfrak{S}_n : D(w) = S\}$$

w	$D(w)$	
123	\emptyset	
213	{1}	$\beta_3(\emptyset) = 1, \quad \beta_3(1) = 2$
312	{1}	$\beta_3(2) = 2, \quad \beta_3(1, 2) = 1$
132	{2}	
231	{2}	
321	{1, 2}	

$\textcolor{red}{u}_S$

Fix n . Let $S \subseteq \{1, \dots, n-1\}$. Let $\textcolor{red}{u}_S = t_1 \cdots t_{n-1}$, where

$$\textcolor{red}{t}_i = \begin{cases} a, & i \notin S \\ b, & i \in S. \end{cases}$$

$\textcolor{red}{u}_S$

Fix n . Let $S \subseteq \{1, \dots, n-1\}$. Let $\textcolor{red}{u}_S = t_1 \cdots t_{n-1}$, where

$$\textcolor{red}{t}_i = \begin{cases} a, & i \notin S \\ b, & i \in S. \end{cases}$$

Example. $n = 8$, $S = \{2, 5, 6\} \subseteq \{1, \dots, 7\}$

$$\textcolor{red}{u}_S = abaabba$$

A noncommutative gen. function

$$\Psi_n(a, b) = \sum_{S \subseteq \{1, \dots, n-1\}} \beta_n(S) u_S.$$

A noncommutative gen. function

$$\Psi_n(a, b) = \sum_{S \subseteq \{1, \dots, n-1\}} \beta_n(S) u_S.$$

Example. Recall

$$\beta_3(\emptyset) = 1, \quad \beta_3(1) = 2, \quad \beta_3(2) = 2, \quad \beta_3(1, 2) = 1$$

Thus

$$\Psi_3(a, b) = aa + 2ab + 2ba + bb$$

A noncommutative gen. function

$$\Psi_n(a, b) = \sum_{S \subseteq \{1, \dots, n-1\}} \beta_n(S) u_S.$$

Example. Recall

$$\beta_3(\emptyset) = 1, \quad \beta_3(1) = 2, \quad \beta_3(2) = 2, \quad \beta_3(1, 2) = 1$$

Thus

$$\Psi_3(a, b) = aa + 2ab + 2ba + bb$$

$$= (a + b)^2 + (ab + ba)$$

The *cd*-index

Theorem. *There exists a noncommutative polynomial $\Phi_n(c, d)$, called the ***cd*-index** of \mathfrak{S}_n , with **nonnegative** integer coefficients, such that*

$$\Psi_n(a, b) = \Phi_n(a + b, ab + ba).$$

The *cd*-index

Theorem. *There exists a noncommutative polynomial $\Phi_n(c, d)$, called the **cd-index** of \mathfrak{S}_n , with **nonnegative** integer coefficients, such that*

$$\Psi_n(a, b) = \Phi_n(a + b, ab + ba).$$

Example. Recall

$$\Psi_3(a, b) = aa + 2ab + 2ba + b^2 = (a + b)^2 + (ab + ba).$$

Therefore

$$\Phi_3(c, d) = c^2 + d.$$

Small values of $\Phi_n(c, d)$

$$\Phi_1 = 1$$

$$\Phi_2 = c$$

$$\Phi_3 = c^2 + d$$

$$\Phi_4 = c^3 + 2cd + 2dc$$

$$\Phi_5 = c^4 + 3c^2d + 5cdc + 3dc^2 + 4d^2$$

$$\begin{aligned}\Phi_6 = & c^5 + 4c^3d + 9c^2dc + 9cdc^2 + 4dc^3 \\ & + 12cd^2 + 10dcd + 12d^2c.\end{aligned}$$

The set S_μ

Recall: $w = a_1 \cdots a_n \in \mathfrak{S}_n$ is **simsun** if for all $1 \leq k \leq n$, the restriction $b_1 \cdots b_k$ of w to $1, 2, \dots, k$ has no **double descents** $b_{i-1} > b_i > b_{i+1}$.

The set S_μ

Recall: $w = a_1 \cdots a_n \in \mathfrak{S}_n$ is **simsun** if for all $1 \leq k \leq n$, the restriction $b_1 \cdots b_k$ of w to $1, 2, \dots, k$ has no **double descents** $b_{i-1} > b_i > b_{i+1}$.

Given a cd -monomial μ , let $c \rightarrow 0$, $d \rightarrow 10$, and remove final 0.

Example.

$$\mu = cd^2c^2d \rightarrow 01010001,$$

the characteristic vector of $S_\mu = \{2, 4, 8\}$.

The coefficients of $\Phi_n(c, d)$

Theorem (Simion-Sundaram) *Let μ be a cd -monomial of degree $n - 1$. The coefficient of μ in $\Phi_n(c, d)$ is the number of simsun permutations $w \in \mathfrak{S}_{n-1}$ with descent set S_μ .*

The case $n = 4$

w	$D(w)$	μ
123	\emptyset	$c^3 \rightarrow 00$
213	1	$dc \rightarrow 10$
312	1	$dc \rightarrow 10$
132	2	$cd \rightarrow 01$
231	2	$cd \rightarrow 01$

$$\Rightarrow \Phi_4(c, d) = c^3 + 2dc + 2cd$$

Expansion of cd-monomials

Let $\textcolor{red}{m} = \mathfrak{m}(c, d)$ be a *cd*-monomial, e.g., c^2d^2cd .
Expand $\mathfrak{m}(a + b, ab + ba)$:

$$(a + b)^2(ab + ba)^2(a + b)(ab + ba) = a^9 + ba^8 + \dots .$$

Expansion of cd-monomials

Let $\textcolor{red}{m} = \text{m}(c, d)$ be a *cd*-monomial, e.g., c^2d^2cd .
Expand $\text{m}(a + b, ab + ba)$:

$$(a + b)^2(ab + ba)^2(a + b)(ab + ba) = a^9 + ba^8 + \dots .$$

Key observation: $\text{m}(a + b, ab + ba)$ is a sum of distinct monomials, always including $ababab\dots$ and $bababa\dots$.

Alternating permutations

The coefficient of $ababab\cdots$ and of $bababa\cdots$ in $\Psi_n(a, b)$ is

$$\beta_n(2, 4, 6, \dots) = \beta_n(1, 3, 5, \dots) = E_n.$$

Alternating permutations

The coefficient of $ababab\cdots$ and of $bababa\cdots$ in $\Psi_n(a, b)$ is

$$\beta_n(2, 4, 6, \dots) = \beta_n(1, 3, 5, \dots) = E_n.$$

Hence:

Theorem. (a) $\Phi_n(1, 1) = E_n$ (*can interpret coefficients combinatorially*).

(b) *For all* $S \subseteq \{1, \dots, n - 1\}$,

$$\beta_n(S) \leq \beta_n(1, 3, 5, \dots) = E_n.$$

An example

$$\Phi_5 = \mathbf{1}c^4 + \mathbf{3}c^2d + \mathbf{5}cdc + \mathbf{3}dc^2 + \mathbf{4}d^2$$

$$1 + 3 + 5 + 3 + 4 = 16 = E_5$$

Comments

NOTE. (b) is due originally to Niven and later de Bruijn (different proofs).

Comments

Note. (b) is due originally to Niven and later de Bruijn (different proofs).

Note. Not hard to show that

$$\beta_n(S) < E_n$$

unless $S = \{1, 3, 5, \dots\}$ or $S = \{2, 4, 6, \dots\}$.

