

THE GENERALIZED KUREPA HYPOTHESIS AT SINGULAR CARDINALS

MOHAMMAD GOLSHANI

ABSTRACT. We discuss the generalized Kurepa hypothesis KH_λ at singular cardinals λ . In particular, we answer questions of Erdős-Hajnal [1] and Todorćevic [6], [7] by showing that GCH does not imply $\text{KH}_{\aleph_\omega}$ nor the existence of a family $\mathcal{F} \subseteq [\aleph_\omega]^{\aleph_0}$ of size $\aleph_{\omega+1}$ such that $\mathcal{F} \upharpoonright X$ has size \aleph_0 for every $X \subseteq S, |X| = \aleph_0$.

1. INTRODUCTION

For an infinite cardinal λ let the generalized Kurepa hypothesis at λ , denoted KH_λ , be the assertion: there exists a family $\mathcal{F} \subseteq P(\lambda)$ such that $|\mathcal{F}| > \lambda$ but $|\mathcal{F} \upharpoonright X| \leq |X|$ for every infinite $X \subseteq \lambda, |X| < \lambda$, where $\mathcal{F} \upharpoonright X = \{t \cap X : t \in \mathcal{F}\}$.

By a theorem of Erdős-Hajnal-Milner [2], if λ is a singular cardinal of uncountable cofinality, $\theta^{\text{cf}(\lambda)} < \lambda$ for all $\theta < \lambda$ and if $\mathcal{F} \subseteq P(\lambda)$ is such that the set $\{\alpha < \lambda : |\mathcal{F} \upharpoonright \alpha| \leq |\alpha|\}$ is stationary in λ , then $|\mathcal{F}| \leq \lambda$. In particular, GCH implies KH_λ fails for all singular cardinals λ of uncountable cofinality. On the other hand, by an unpublished result of Prikry [5], KH_λ holds in L , the Gödel's constructible universe, for singular cardinals of countable cofinality (see [7]). Later, Todorćevic [6], [7] improved Prikry's theorem by showing that if λ is a singular cardinal of countable cofinality, then \square_λ implies KH_λ . The following question is asked in [6] and [7].

Question 1.1. *Does GCH imply $\text{KH}_{\aleph_\omega}$?*

The question is also related to the following question of Erdős-Hajnal [1] (question 19/E)

Question 1.2. *Assume GCH. Let $|S| = \aleph_\omega$. Does there exist a family $\mathcal{F}, |\mathcal{F}| = \aleph_{\omega+1}, \mathcal{F} \subseteq [S]^{\aleph_0}$ such that $\mathcal{F} \upharpoonright X$ has size \aleph_0 for every $X \subseteq S, |X| = \aleph_0$?*

The author's research has been supported by a grant from IPM (No. 97030417). He also thanks Menachem Magidor for sharing his ideas during the 14th International Workshop in Set Theory at Luminy, in particular, the results of this paper are obtained from discussions with him.

We show that, relative to the existence of large cardinals, both of the above questions can consistently be false, and so they are independent of ZFC.

2. KH_λ FAILS ABOVE A SUPERCOMPACT CARDINAL

In this section we prove the following.

Theorem 2.1. *Suppose κ is a supercompact cardinal and $\lambda \geq \kappa$. Then KH_λ fails.*

Proof. Let $\mathcal{F} \subseteq P(\lambda)$ be of size $\geq \lambda^+$. Let $j : V \rightarrow M$ be a λ^+ -supercompactness embedding with $\text{crit}(j) = \kappa$. Also let U be the normal measure on $P_\kappa(\lambda)$ derived from j , i.e.,

$$U = \{X \subseteq P_\kappa(\lambda) : j[\lambda] \in j(X)\}.$$

We have

- $M \models "j(\mathcal{F}) \subseteq P(j(\lambda))$ is of size $\geq j(\lambda)^+$ ".
- $j''[\lambda] \in M$ and $M \models "|j''[\lambda]| = \lambda < j(\lambda)"$.
- $\mathcal{F} \in M$
- $a \neq b \in \mathcal{F} \implies j(a) \cap j''[\lambda] \neq j(b) \cap j''[\lambda]$.

In particular,

$$M \models "|j(\mathcal{F}) \restriction j''[\lambda]| \geq |\mathcal{F}| \geq \lambda^+."$$

This implies that

$$\{x \in P_\kappa(\lambda) : |\mathcal{F} \restriction x| \geq |x|^+\} \in U.$$

In particular, \mathcal{F} is not a KH_λ -family. □

Remark 2.2. *The above result is optimal in the sense that we can not in general find a set $x \subseteq \lambda$ of size in the interval $[\kappa, \lambda)$ such that $|\mathcal{F} \restriction x| \geq |x|^+$. To see this assume κ is supercompact and Laver indestructible. Then one can easily define a κ -directed closed forcing notion which adds a family $\mathcal{F} \subseteq P(\lambda)$ such that $|\mathcal{F}| \geq \lambda^+$, but $|\mathcal{F} \restriction x| \leq |x|$ for any set x with $\kappa \leq |x| < \lambda$.*

3. THE CHANG'S CONJECTURE AND $\text{KH}_{\aleph_\omega}$

In this section we prove our main theorem by showing a consistent negative answer to the questions of Erdős-Hajnal and Todorćevic. Recall from [4] that “GCH + the Chang’s conjecture $(\aleph_{\omega+1}, \aleph_\omega) \rightarrow (\aleph_1, \aleph_0)$ ” is consistent, relative to the existence of a 2-huge cardinal. See also [3], where the large cardinal assumption is reduced to the existence of a $(+\omega + 1)$ -subcompact cardinal κ .

Theorem 3.1. *Assume GCH + Chang’s conjecture $(\aleph_{\omega+1}, \aleph_\omega) \rightarrow (\aleph_1, \aleph_0)$. Then $\text{KH}_{\aleph_\omega}$ fails. Also, there does not exist a family $\mathcal{F} \subseteq [\aleph_\omega]^{\aleph_0}$, $|\mathcal{F}| \geq \aleph_{\omega+1}$ such that $\mathcal{F} \upharpoonright X$ has size \aleph_0 for every $X \subseteq S$, $|X| = \aleph_0$.*

Proof. Suppose towards contradiction that there exists a family \mathcal{F} which witnesses $\text{KH}_{\aleph_\omega}$. Fix a bijection $f : H_{\aleph_{\omega+1}} \leftrightarrow \mathcal{F}$. Consider the structure

$$\mathcal{A} = (H_{\aleph_{\omega+1}}, \in, \mathcal{F}, \aleph_\omega, f).$$

Let $\mathcal{B} = (B, \in, \mathcal{G}, A, g) \prec \mathcal{A}$ be such that $|B| = \aleph_1$ and $|A| = \aleph_0$.

Note that $\mathcal{A} \models \forall t \in \mathcal{F}, t \subseteq \aleph_\omega$, and hence $\mathcal{B} \models \forall t \in \mathcal{G}, t \subseteq A$, in particular, $\mathcal{G} \subseteq \mathcal{F} \upharpoonright A$. On the other hand $g : B \leftrightarrow \mathcal{G}$ is a bijection, hence we have

$$|\mathcal{F} \upharpoonright A| \geq |\mathcal{G}| = |B| = \aleph_1 > \aleph_0.$$

We get a contradiction and the result follows.

Similar argument shows that there can not be a family $\mathcal{F} \subseteq [\aleph_\omega]^{\aleph_0}$ as stated above. \square

REFERENCES

- [1] Erdős, P.; Hajnal, A., Unsolved problems in set theory. 1971 Axiomatic Set Theory (Proc. Sympos. Pure Math., Vol. XIII, Part I, Univ. California, Los Angeles, Calif., 1967) pp. 17-48 Amer. Math. Soc., Providence, R.I.
- [2] Erdős, P.; Hajnal, A.; Milner, E. C., On sets of almost disjoint subsets of a set. Acta Math. Acad. Sci. Hungar 19 1968 209-218.
- [3] Hayut, Yair, Magidor-Malitz reflection. Arch. Math. Logic 56 (2017), no. 3–4, 253-272.
- [4] Levinski, Jean-Pierre; Magidor, Menachem; Shelah, Saharon, Chang’s conjecture for \aleph_ω . Israel J. Math. 69 (1990), no. 2, 161-172.
- [5] Prikry, Karel, Kurepa’s hypothesis for singular cardinals, unpublished note.

- [6] Todorcevic, Stevo, Trees and linearly ordered sets. Handbook of set-theoretic topology, 235-293, North-Holland, Amsterdam, 1984.
- [7] Todorcevic, Stevo, Aronszajn trees and partitions. Israel J. Math. 52 (1985), no. 1-2, 53-58.

Mohammad Golshani, School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O. Box: 19395-5746, Tehran-Iran.

E-mail address: golshani.m@gmail.com

URL: <http://math.ipm.ac.ir/golshani/>